

Generalized Bivariate Fibonacci Polynomials

Mario Catalani

Department of Economics, University of Torino
Via Po 53, 10124 Torino, Italy
mario.catalani@unito.it

Abstract

We define generalized bivariate polynomials, from which specifying initial conditions the bivariate Fibonacci and Lucas polynomials are obtained. Using essentially a matrix approach we derive identities and inequalities that in most cases generalize known results.

1 Antefacts

The generalized bivariate Fibonacci polynomial may be defined as

$$H_n(x, y) = xH_{n-1}(x, y) + yH_{n-2}(x, y), \quad H_0(x, y) = a_0, H_1(x, y) = a_1.$$

We assume $y \neq 0$, as well as $x^2 + 4y \neq 0$.

If we set $a_0 = 0, a_1 = 1$ we obtain the bivariate Fibonacci polynomials $F_n(x, y)$; with $a_0 = 2, a_1 = x$ we get the bivariate Lucas polynomials $L_n(x, y)$. The univariate polynomials are obtained setting $y = 1$. The characteristic polynomial is

$$f(t) = t^2 - tx - y,$$

and the roots are (see for example [3])

$$\alpha \equiv \alpha(x, y) = \frac{x + \sqrt{x^2 + 4y}}{2}, \quad \beta \equiv \beta(x, y) = \frac{x - \sqrt{x^2 + 4y}}{2}.$$

Note that $\alpha + \beta = x, \alpha\beta = -y$. Furthermore $\alpha > \beta$; if $y > 0$ then $\alpha > 0$ and $\beta < 0$ and if $x > 0$ then $\alpha > |\beta|$, while if $x < 0$ then $\alpha < |\beta|$; if $y < 0$ then α and β have the same sign: if $x > 0$ they are both positive, while if $x < 0$ they are both negative.

The ordinary generating function $g(t)$ is

$$g(t) = \frac{a_0 + (a_1 - a_0x)t}{1 - xt - yt^2},$$

which can be obtained through the usual trick: write

$$g(t) = H_0 + H_1t + H_2t^2 + \dots$$

then obtain $xtg(t)$ and $yt^2g(t)$, subtract them from $g(t)$, use the recurrence to get rid of all summands except the first two and impose the initial conditions. Using the Rational Expansion Theorem (see [5]) we get the Binet's form

$$H_n(x, y) = (a_1 - \beta a_0) \frac{\alpha^n}{\alpha - \beta} - (a_1 - \alpha a_0) \frac{\beta^n}{\alpha - \beta}. \quad (1)$$

Hence

$$F_n(x, y) = \frac{\alpha^n}{\alpha - \beta} - \frac{\beta^n}{\alpha - \beta},$$

$$L_n(x, y) = \alpha^n + \beta^n.$$

In the sequel we will write simply H_n , F_n and L_n instead of $H_n(x, y)$, $F_n(x, y)$ and $L_n(x, y)$.

The sequence can be extended to negative subscripts by defining

$$H_{-n} = -\frac{x}{y}H_{-(n-1)} + \frac{1}{y}H_{-(n-2)}.$$

In this case the characteristic polynomial is the reflected polynomial (see [5]) of $f(t)$: hence the roots are $\frac{1}{\alpha}$ and $\frac{1}{\beta}$. Define the matrix

$$\mathbf{A} = \begin{bmatrix} x & 1 \\ y & 0 \end{bmatrix}. \quad (2)$$

Note that $|\mathbf{A}| = -y$.

By easy induction

$$\mathbf{A}^n = \begin{bmatrix} F_{n+1} & F_n \\ yF_n & yF_{n-1} \end{bmatrix}. \quad (3)$$

From the above equation taking determinants

$$F_{n-1}F_{n+1} - F_n^2 = (-1)^n y^{n-1}. \quad (4)$$

Also from $\mathbf{A}^{2n} = \mathbf{A}^n \mathbf{A}^n$ it follows

$$F_{n+1}^2 + yF_n^2 = F_{2n+1}. \quad (5)$$

The eigenvalues of $\mathbf{A}$ are α and β : from this

$$\text{tr}(\mathbf{A}^n) = F_{n+1} + yF_{n-1} = \alpha^n + \beta^n = L_n. \quad (6)$$

Also it is immediate to get

$$F_n L_n = F_{2n}.$$

A wealth of inequalities can be obtained from Equation 3 using the classical matrix inequality on the extremes of quadratic forms

$$\beta^n(z_1^2 + z_2^2) \leq z_1^2 F_{n+1} + z_1 z_2 (1 + y) F_n + z_2^2 y F_{n-1} \leq \alpha^n(z_1^2 + z_2^2), \quad (7)$$

for all real z_1, z_2 .

Now define the matrix

$$\begin{aligned} \mathbf{C} &= \begin{bmatrix} a_1 x + a_0 y & a_1 \\ a_1 y & a_0 y \end{bmatrix} \\ &= a_0 y \mathbf{I} + a_1 \mathbf{A}. \end{aligned}$$

Again by easy induction we get

$$\mathbf{C} \mathbf{A}^n = \begin{bmatrix} H_{n+2} & H_{n+1} \\ y H_{n+1} & y H_n \end{bmatrix}. \quad (8)$$

Taking $a_0 = 2, a_1 = x$ we get the matrix

$$\begin{aligned} \mathbf{B} &= \begin{bmatrix} x^2 + 2y & x \\ xy & 2y \end{bmatrix} \\ &= 2y \mathbf{I} + x \mathbf{A}, \end{aligned}$$

and

$$\mathbf{B} \mathbf{A}^n = \begin{bmatrix} L_{n+2} & L_{n+1} \\ y L_{n+1} & y L_n \end{bmatrix}. \quad (9)$$

Because

$$\mathbf{C} \mathbf{A}^n = (a_0 y \mathbf{I} + a_1 \mathbf{A}) \mathbf{A}^n = a_0 y \mathbf{A}^n + a_1 \mathbf{A}^{n+1},$$

we obtain the identity

$$H_{n+1} = a_0 y F_n + a_1 F_{n+1},$$

which specializes to

$$L_{n+1} = 2y F_n + x F_{n+1}.$$

Analogous results we get for H_{-n} using the inverse of $\mathbf{A}$

$$\mathbf{A}^{-1} = \begin{bmatrix} 0 & \frac{1}{y} \\ 1 & -\frac{x}{y} \end{bmatrix}.$$

In particular we get

$$H_{-n} = \frac{(-1)^n}{y} [a_0 F_{n+1} - a_1 F_n].$$

Because $|\mathbf{C}| = a_0 a_1 x y + a_0^2 y^2 - a_1^2 y$ taking determinants in Equation 8 we get

$$H_n H_{n+2} - H_{n+1}^2 = (-1)^n y^{n-1} (a_0 a_1 x y + a_0^2 y^2 - a_1^2 y), \quad (10)$$

which can be termed the *Simpson* formula.

Using identity 1.60 in [4], that is

$$\sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} (xy)^k (x+y)^{n-2k} = \frac{x^{n+1} - y^{n+1}}{x - y},$$

writing α for x and β for y we have

$$\begin{aligned} \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k \binom{n-k}{k} (\alpha\beta)^k (\alpha + \beta)^{n-2k} &= \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} \\ &= F_{n+1}, \end{aligned}$$

which means

$$\sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-k}{k} y^k x^{n-2k} = F_{n+1}.$$

2 Inversion formulas

Now we will establish the following formula for γ complex and $i^2 = -1$

$$F_n(x, -y) = (i\gamma)^{-(n-1)} F_n(i\gamma x, \gamma^2 y). \quad (11)$$

We have

$$F_n(x, -y) = \frac{\alpha_1^n - \beta_1^n}{\alpha_1 - \beta_1},$$

where

$$\alpha_1 = \frac{x + \sqrt{x^2 - 4y}}{2}.$$

Analogously for β_1 . Also

$$F_n(i\gamma x, \gamma^2 y) = \frac{\alpha_2^n - \beta_2^n}{\alpha_2 - \beta_2},$$

where

$$\begin{aligned} \alpha_2 &= \frac{i\gamma x + \sqrt{-\gamma^2 x^2 + 4\gamma^2 y}}{2} \\ &= \frac{i\gamma x + i\gamma \sqrt{x^2 - 4y}}{2} \\ &= i\gamma \frac{x + \sqrt{x^2 - 4y}}{2} \\ &= i\gamma \alpha_1. \end{aligned}$$

Analogously for β_2 . Then

$$\begin{aligned} F_n(i\gamma x, \gamma^2 y) &= \frac{(i\gamma)^n \alpha_1^n - (i\gamma)^n \beta_1^n}{i\gamma \alpha_1 - i\gamma \beta_1} \\ &= (i\gamma)^{n-1} F_n(x, -y), \end{aligned}$$

whence the desired formula. Some consequences. If we set $\gamma = i$ we get

$$F_n(x, -y) = (-1)^{n-1} F_n(-x, -y). \quad (12)$$

Also from

$$F_n(x, y) = (i\gamma)^{-(n-1)} F_n(i\gamma x, -\gamma^2 y)$$

we get with $\gamma = \frac{1}{\sqrt{-y}} = \frac{1}{i\sqrt{y}}$

$$F_n(x, y) = (\sqrt{y})^{n-1} F_n\left(\frac{x}{\sqrt{y}}, 1\right). \quad (13)$$

In a similar fashion we obtain

$$L_n(x, -y) = (i\gamma)^{-n} L_n(i\gamma x, \gamma^2 y), \quad (14)$$

$$L_n(x, y) = (\sqrt{y})^n L_n\left(\frac{x}{\sqrt{y}}, 1\right),$$

and

$$L_n(-x, y) = (-1)^n L_n(x, y).$$

It follows

$$F_{2n+1}(-x, y) = F_{2n+1}(x, y), \quad F_{2n}(-x, y) = -F_{2n}(x, y),$$

$$L_{2n+1}(-x, y) = -L_{2n+1}(x, y), \quad L_{2n}(-x, y) = F_{2n}(x, y).$$

From the Binet's forms we get

$$F_n(ax, by) = a^{n-1} F_n\left(x, \frac{b}{a^2} y\right)$$

$$L_n(ax, by) = a^n L_n\left(x, \frac{b}{a^2} y\right)$$

from which we get

$$F_n(x, y) = x^{n-1} F_n\left(1, \frac{y}{x^2}\right),$$

$$L_n(x, y) = x^n L_n\left(1, \frac{y}{x^2}\right).$$

3 A complex representation

Consider the matrix

$$\mathbf{B} = \frac{1}{\sqrt{x+2i}} \begin{bmatrix} x+i & 1 \\ 1 & i \end{bmatrix},$$

where $i = \sqrt{-1}$. Then

$$\mathbf{B}^2 = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix} = \mathbf{A}.$$

This matrix is obtained by Equation 2 setting $y = 1$ and so is the matrix generating the univariate Fibonacci polynomials. Then

$$\mathbf{A}^n = \begin{bmatrix} F_{n+1}(x, 1) & F_n(x, 1) \\ F_n(x, 1) & F_{n-1}(x, 1) \end{bmatrix}.$$

Using the method exposed in [7] to $\mathbf{B}^{2n} = \mathbf{A}^n$ (which consists in expressing the elements of the n -th power of a 2×2 matrix in terms of the elements of the matrix) we get

$$F_n(x, 1) = \sum_{r=0}^{n-1} \binom{2n-1-r}{r} (x+2i)^{n-r-1} (-i)^r. \quad (15)$$

Using Equation 13, after replacing in the previous equation x by $\frac{x}{\sqrt{y}}$, we get

$$F_n(x, y) = \sum_{r=0}^{n-1} \binom{2n-1-r}{r} (x+2\sqrt{y}i)^{n-r-1} (-\sqrt{y}i)^r. \quad (16)$$

4 Summations

Since

$$\sum_{k=0}^n \alpha^k = \frac{1 - \alpha^{n+1}}{1 - \alpha},$$

(analogously for β) and $(1 - \alpha)(1 - \beta) = 1 - x - y$, we have

$$\begin{aligned} \sum_{k=0}^n H_k &= \frac{a_1 - a_0\beta}{\alpha - \beta} \sum_{k=0}^n \alpha^k - \frac{a_1 - a_0\alpha}{\alpha - \beta} \sum_{k=0}^n \beta^k \\ &= \frac{a_0(1 - x) + a_1 - yH_n - H_{n-1}}{1 - x - y}, \end{aligned}$$

which can be rewritten as

$$(1 - x - y) \sum_{k=0}^n H_k = a_0(1 - x) + a_1 - yH_n - H_{n-1}.$$

Specializing we get

$$(1 - x - y) \sum_{k=0}^n F_k = 1 - yF_n - F_{n+1},$$

$$(1 - x - y) \sum_{k=0}^n L_k = 2 - x - yL_n - L_{n+1}.$$

Since

$$\sum_{k=0}^n \alpha^{2k} = \frac{1 - \alpha^{2n+2}}{1 - \alpha^2},$$

(analogously for β), $(1-\alpha^2)(1-\beta^2) = 1-x^2-2y+y^2$, and $\alpha^2-\beta^2 = x(\alpha-\beta)$ we have

$$(1-2y-x^2+y^2) \sum_{k=0}^n F_{2k} = x + y^2 F_{2n} - F_{2n+2},$$

$$(1-2y-x^2+y^2) \sum_{k=0}^n L_{2k} = 2-x^2-2y+y^2 L_{2n} - L_{2n+2}.$$

Just a little bit more involved is the following inequality. Consider

$$\begin{aligned} L_n^m &= (\alpha^n + \beta^n)^m \\ &= \sum_{k=0}^m \binom{m}{k} \alpha^{nk} \beta^{n(m-k)}. \end{aligned}$$

Now upon expanding the summation and grouping the first and the last element, the second and the next-to-last element and so on we get the following formulas depending on the parity of m

$$L_n^m = \sum_{k=0}^{\lfloor \frac{m}{2} \rfloor} \binom{m}{k} (-y)^{kn} L_{n(m-2k)} \quad \text{if } m \text{ is odd,}$$

$$L_n^m = \sum_{k=0}^{\frac{m}{2}-1} \binom{m}{k} (-y)^{kn} L_{n(m-2k)} + \binom{m}{\frac{m}{2}} (-y)^{\frac{mn}{2}} \quad \text{if } m \text{ is even.}$$

If we define

$$\varepsilon_1 = \left\lfloor \frac{m}{2} \right\rfloor + \frac{1}{2} [-1 + (-1)^{m+1}], \quad \varepsilon_2 = \frac{1}{2} [(-1)^m + 1],$$

we can group the two identities into the following one

$$L_n^m = \sum_{k=0}^{\varepsilon_1} \binom{m}{k} (-y)^{kn} L_{n(m-2k)} + \varepsilon_2 \binom{m}{\frac{m}{2}} (-y)^{\frac{mn}{2}}.$$

Other consequences are

$$\sum_{k=0}^n \binom{n}{k} x^k y^{n-k} F_k = F_{2n},$$

$$\sum_{k=0}^n \binom{n}{k} x^k y^{n-k} L_k = L_{2n}.$$

Indeed, for instance the second one, we have

$$\begin{aligned}\sum_{k=0}^n \binom{n}{k} x^k y^{n-k} L_k &= \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} \alpha^k + \sum_{k=0}^n \binom{n}{k} x^k y^{n-k} \beta^k \\ &= (y + x\alpha)^n + (y + x\beta)^n,\end{aligned}$$

and $y + x\alpha = \alpha^2$, $y + x\beta = \beta^2$.

5 Eigenvectors and eigenvalues

Now $\forall k, n$

$$\begin{aligned}\mathbf{A}^k \mathbf{B} \mathbf{A}^n &= \mathbf{A}^k (y\mathbf{I} + \mathbf{A}^2) \mathbf{A}^n \\ &= y\mathbf{A}^{n+k} + \mathbf{A}^{n+k+2} \\ &= (y\mathbf{I} + \mathbf{A}^2) \mathbf{A}^{n+k} \\ &= \mathbf{B} \mathbf{A}^{n+k}.\end{aligned}$$

This means that $\forall k, n$ the matrices $\mathbf{A}^k$ and $\mathbf{B} \mathbf{A}^n$ commute: then (see [1], [2], [8]) the matrices possess the same eigenvectors.

The eigenvectors (not normalized) of $\mathbf{A}$ are

$$\mathbf{z}_1 = \begin{bmatrix} \alpha \\ y \\ 1 \end{bmatrix}, \quad \text{and} \quad \mathbf{z}_2 = \begin{bmatrix} \beta \\ y \\ 1 \end{bmatrix}.$$

For the eigenvalues of $\mathbf{B} \mathbf{A}^n$ since

$$\mathbf{B} \mathbf{A}^n = y\mathbf{A}^n + \mathbf{A}^{n+2}$$

we have

$$\begin{cases} \lambda_1 = y\alpha^n + \alpha^{n+2}, \\ \lambda_2 = y\beta^n + \beta^{n+2}. \end{cases}$$

Consider $(\mathbf{B} \mathbf{A}^n)^2 = \mathbf{B} \mathbf{A}^n \mathbf{B} \mathbf{A}^n$. Since $\mathbf{A}^k$ and $\mathbf{B} \mathbf{A}^n$ commute, this means that we can write $(\mathbf{B} \mathbf{A}^n)^2 = \mathbf{B} \mathbf{B} \mathbf{A}^{2n}$, that is

$$\begin{bmatrix} 2y + x^2 & x \\ xy & 2y \end{bmatrix} \begin{bmatrix} L_{2n+2} & L_{2n+1} \\ yL_{2n+1} & yL_{2n} \end{bmatrix}.$$

Using repeatedly the recurrence for L_n we get

$$L_{n+2}^2 + yL_{n+1}^2 = L_{2n+4} + yL_{2n+2}. \quad (17)$$

Another immediate consequence is the following. Since

$$\mathbf{A}^n \mathbf{B} \mathbf{A}^n = \begin{bmatrix} F_{n+1}L_{n+2} + yF_nL_{n+1} & F_{n+1}L_{n+1} + yF_nL_n \\ yF_nL_{n+2} + y^2F_{n-1}L_{n+1} & yF_nL_{n+1} + y^2F_{n-1}L_n \end{bmatrix},$$

and $\mathbf{A}^n \mathbf{B} \mathbf{A}^n = \mathbf{B} \mathbf{A}^{2n}$ it follows equating the elements in position $(2, 2)$

$$L_{2n} = F_nL_{n+1} + yF_{n-1}L_n. \quad (18)$$

Because of the *Schur inequality* (see [1], [6]), that is: the sum of the squared elements is greater or equal to the sum of the squared eigenvalues, we have the following inequality

$$\begin{aligned} F_{n+1}^2 + (1+y^2)F_n^2 + y^2F_{n-1}^2 &\geq \alpha^{2n} + \beta^{2n} \\ &= L_{2n}. \end{aligned}$$

Note that the equality sign holds if $y = 1$.

In the same way we get

$$\begin{aligned} y^2L_n^2 + (1+y^2)L_{n+1}^2 + L_{n+2}^2 &\geq (y\alpha^n + \alpha^{n+2})^2 + (y\beta^n + \beta^{n+2})^2 \\ &= y^2L_{2n} + 2yL_{2n+2} + L_{2n+4}, \end{aligned}$$

and the equality sign holds if $y = 1$.

Consider

$$\begin{aligned} \mathbf{B}^2 \mathbf{A}^n &= y^2 \mathbf{A}^n + \mathbf{A}^{n+4} + 2y \mathbf{A}^{n+2} \\ &= y \mathbf{B} \mathbf{A}^n + \mathbf{B} \mathbf{A}^{n+2} \\ &= \begin{bmatrix} yL_{n+2} + L_{n+4} & yL_{n+1} + L_{n+3} \\ y^2L_{n+1} + yL_{n+3} & y^2L_n + yL_{n+2} \end{bmatrix}. \end{aligned}$$

Taking the determinants we get

$$\begin{aligned} (-1)^n y^n (x^2 y + 4y^2)^2 &= y^3 L_n L_{n+2} + y^2 L_{n+2}^2 + y^2 L_n L_{n+4} + y L_{n+2} L_{n+4} \\ &\quad - y^3 L_{n+1}^2 - y L_{n+3}^2 - 2y^2 L_{n+1} L_{n+3}. \end{aligned}$$

Using repeatedly identity 10 we obtain

$$L_n L_{n+4} - L_{n+1} L_{n+3} = (-1)^n (x^4 y^n + 5x^2 y^{n+1} + 4y^{n+2}). \quad (19)$$

References

- [1] M. Catalani (1996), *Spazi Lineari*, Iuculano Editore, Pavia.
- [2] F.R. Gantmacher (1977), *The Theory of Matrices*, Chelsea Publishing Company, New York.
- [3] N. Gauthier and J.R. Gosselin (2002), "Problem H-587." *The Fibonacci Quarterly* **40.4**: 380.
- [4] H.W. Gould (1972), *Combinatorial Identities*, Morgantown, W. Va.
- [5] R. Graham, D. Knuth and O. Patashnik (1998), *Concrete Mathematics*, Addison-Wesley, Reading, MA.
- [6] M. Marcus and H. Minc (1969), *A Survey of Matrix Theory and Matrix Inequalities*, Dover Publications, New York.
- [7] J. Mc Laughlin (2004), "Combinatorial Identities Deriving from the n -th Power of a 2×2 Matrix", published electronically at <http://emp.trincoll.edu/~jmclaugh/2x2atrixpowerglaintegers.pdf>
- [8] C.R. Rao and S.K. Mitra (1971), *Generalized Inverse of Matrices and its Applications*, Wiley and Sons, New York.